



Manufactured in U.S.A. by:

SCI-PHARM®

Sci-Pharm Catalog No. 85-30
TEETH WHITENING SYSTEM
WITH CALCIUM

TECHNICAL BULLETIN
Instructions

Fast & Gentle™

Teeth Whitening System With Calcium

**FOR APPLICATION BY A PROFESSIONAL OR
UNDER PROFESSIONAL SUPERVISION**

KIT CONTAINS:

- 14g Teeth Whitener in two automix syringes
- 4 Static mixers (tips)
- 2 Tray forming sheets
- Instructions



OUTSTANDING FEATURES OF THE MATERIAL

- **Fast Acting; short application time**
- **Low potential for tissue irritation**
- **User friendly, easy to handle automix dispensing system**
- **Short treatment time**
- **Requires no special equipment**
- **Contains calcium to prevent tooth decalcification during treatment**

STORAGE AND SHELF LIFE

The shelf life of the material is six months when stored at temperatures not exceeding 73°F(23°C). Constant storage under refrigeration will extend the shelf life of the material to one year.

Produced under one or more of the following U.S. Patents: 6,365,134, 6,447,757, 6,488,913, 6,696,043, 7,067,115, Other patents pending

CAUTION

Federal law restricts this device to sale
by or on the order of a dentist.

SPECIAL COMMENTS

This teeth whitening system is most efficient if treatments are done consecutively, within a one to two week time period.

After placing the mixing tip onto the syringe, do not remove the tip until all the material has been used. Between uses, the material may form a seal over the tip of the syringe. Immediately before the next treatment, dispense and discard a small, pea sized amount of the material. The syringe is then ready to fill the trays. If precipitation of the material in the tip between uses makes the dispensing difficult, remove the tip and clean it in hot water to loosen the material, and then expel the loosened material.

Each of the dual barrel syringes has enough material for about six applications.

INSTRUCTIONS

1. Prior to first treatment, procure the trays and adjust to achieve a good fit.
2. Place the static mixing tip onto the dual barrel syringe and twist clockwise to lock into place. After each application, leave the mixing tip on the syringe. Do not retract the plunger, even slightly, as this can cause inadvertent activation of the material in the syringe.

Note: After each application, a small amount of the mixed material will remain in the static mixer. Do not remove the material from the tip; this will result in desirable sealing of the orifice.

3. Dispense the mixed contents of the two barrels onto that portion of the tray that will cover the labial surfaces of the teeth to be whitened.
4. Place tray into mouth. Remove any excess of the material with tissue.
5. After 30 to 45 minutes, remove tray and rinse. If there is no discomfort, the next treatment time may be extended up to sixty minutes. Prolonging the time of individual treatments will reduce the number of applications necessary to achieve a desirable effect.
6. Repeat treatments once or twice a day or as directed by a dentist. Do not eat, drink, or smoke while trays are in place.

Warning: This product should not be used on children under 12, on pregnant or nursing women, or on individuals allergic to any of the ingredients listed.

INGREDIENTS

Glycerin, Carbamide Peroxide, Propylene Glycol, Water, Calcium Peroxide, Polyethylether, Calcium Carbonate, Silica, Sodium Pyrophosphate, Titanium Dioxide, 1-Hydroxyethylidene-phosphonic Acid, Polyacrylic Acid.

For technical information, call or write:



CE
0510

EU Authorized Representative:
QUALITY FIRST INTERNATIONAL LIMITED
20 Eversley Road, Bexhill-on-Sea, East Sussex TN40 1HE, UK
Voice: + 44 20 8 522 1937 • Fax: + 44 20 8 522 1937
E-mail: Hard6162@aol.com

SCIENTIFIC PHARMACEUTICALS, INC.
3221 PRODUCER WAY • POMONA, CALIFORNIA USA 91768
PHONE: (800) 634-3047 • (909) 595-9922
FAX: (909) 595-0331 • E-MAIL: scipharm@msn.com
WEBSITE: www.scipharm.com